## DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

T. B. C.: DS - 2015

Seriál No.

1784

**Test Booklet Series** 



## **TEST BOOKLET**

**DENTAL SURGEON** 

Time Allowed : 3 Hours

(Maximum Marks : 200)

## : INSTRUCTIONS TO CANDIDATES :

- 1. IMMEDIATELY AFTER THE COMMENCEMENT OF THE EXAMINATION, YOU SHOULD CHECK THAT THIS TEST BOOKLET **DOES NOT** HAVE ANY UNPRINTED OR TORN OR MISSING PAGES OR ITEMS ETC. IF SO, GET IT REPLACED BY A COMPLETE TEST BOOKLET OF THE SAME SERIES ISSUED TO YOU.
- 2. ENCODE CLEARLY THE TEST BOOKLET SERIES **A**, **B**, **C** OR **D**, AS THE CASE MAY BE, IN THE APPROPRIATE PLACE IN THE ANSWER SHEET USING BALL POINT PEN (BLUE OR BLACK).
- 3. You have to enter your Roll No. on the Test Booklet in the Box provided alongside. DO NOT write anything else on the Test Booklet.
- 4. This Test Booklet contains 200 items (questions). Each item (question) comprises four responses (answers). You have to select the correct response (answer) which you want to mark (darken) on the Answer Sheet. In case, you feel that there is more than one correct response (answer), you should mark (darken) the response (answer) which you consider the best. In any case, choose ONLY ONE response (answer) for each item (question).
- You have to mark (darken) all your responses (answers) ONLY on the separate Answer Sheet provided, by using BALL POINT PEN (BLUE OR BLACK). See instructions in the Answer Sheet.
- 6. All items (questions) carry equal marks. All items (questions) are compulsory. Your total marks will depend only on the number of correct responses (answers) marked by you in the Answer Sheet. There will be no negative marking for wrong answer.
- 7. Before you proceed to mark (darken) in the Answer Sheet the responses to various items (questions) in the Test Booklet, you have to fill in some particulars in the Answer Sheet as per the instructions sent to you with your **Admission Certificate**.
- 8. After you have completed filling in all your responses (answers) on the Answer Sheet and after conclusion of the examination, you should hand over to the Invigilator the *Answer Sheet* issued to you. You are allowed to take with you the candidate's copy/second page of the Answer Sheet along with the *Test Booklet* after completion of the examination for your reference.

DO NOT OPEN THIS TEST BOOKLET UNTIL YOU ARE ASKED TO DO SO

- "Enter bolus-exit balance" was stated by:
   (A) Harold Ortman
   (B) Sheppard
   (C) Stewart
   (D) Hudson
- 2. The estimated width of maxillary central incisor is:
  - (A) 1/16<sup>th</sup> bizygomatic width
  - (B) 1/16<sup>th</sup> interpupillary distance
  - (C) 1/8<sup>th</sup> bizygomatic width
  - (D) None of the above
- 3. In arcon articulators the condylar element is located in :
  - (A) Upper member
  - (B) Lower member
  - (C) No condyle
  - (D) Fixed condyle
- 4. The quality that distinguishes one from the other is:
  - (A) Hue
  - (B) Chroma
  - (C) Value
  - (D) Saturation
- 5. In laminate technique of impression making :
  - (A) Alginate and agar are used
  - (B) Only alginate is used
  - (C) Elastomers are used
  - (D) Only agar

- 6. Central incisors are generally anterior to the line bisecting incisive pappilae by:
  - (A) 8-10 mm
  - (B) 14-15 mm
  - (C) 2-6 mm
  - (D) 11-20 mm
- 7. Advantages of an over denture:
  - (A) Preservation of ridge
  - (B) Support
  - (C) Retention
  - (D) All of the above
- 8. Facebow transfer records:
  - (A) Orientation relation
  - (B) Special relation
  - (C) Vertical relation
  - (D) Tentative centric relation
- Identify the correct hanau formula which gives the lateral condylar guidance angle:
  - (A) L = H/8 + 12
  - (B) L = 8/H + 12
  - (C) L = H/8 \* 12
  - (D) L = H/8 12
- 10. Instrument used to locate pterygomaxillary notch:
  - (A) Boby gauge
  - (B) Divider
  - (C) Screw gauge
  - (D) T-burnisher

- 11. The distal border of mandibular 16. denture should cover: (A) Ant, <sup>1</sup>/<sub>3</sub>rd of retromolar pad Methyl red (B) Glucose Ant. ard of retromolar pad (C) (D) (C) Should not cover the retromolar pad (D) Hamullar notch (A) London In acrylic resin, hydroquinone act as: (B) New York (A) Activator (C) Grand Rapids (B) Initiator (C) Inhibitor 18. (D) Plasticizer True Positive (A) 13. Alginate is made dust free by the (B) **True Negative** addition of: **False Positive** (C) Glycerine (A) (D) False Negative (B) Calcium sulphate (C) Sodium phosphate Calcium chloride 14. Rest seat can be prepared: Cartogram (B) (A) Only on tooth surface (C) Histogram (B) In amalgam restoration Pie diagram (D) (C) In cast gold restoration All of the above
- 15. Heal raising movement of a RPD is prevented by:
  - (A) Occlusal rest
  - Minor connector (B)
  - (C) Direct retainer
  - Indirect retainer

- Indicator in Snyder Test is:
  - Diazoresorcinol
  - Bromocresol green
- 17. First artificial water fluoridation of water supply was done in:
  - (D) Ontario, Canada
- Specificity of a test denotes:

- 19. The coverage of the geographic area through dental clinics can be depicted by the following way:
  - (A) Frequency polygon
- 20. An experiment in which the experimental subjects cannot distinguish the experimental conditions is called as:
  - (A) Two Tailed Testing
  - Single Blind experiment (B)
  - **Double Blind experiment** (C)
  - Retrospective experiment (D)

21.	Pit a	nd fissure sealant is a form of :	26.	Circ	umoral pigmentation is seen in :
	(A)	Primary prevention		(A)	Arsenic poisoning
	(B)	Secondary prevention		(B)	Lead poisoning
	(C)	Tertiary prevention		(C)	Peutz-Jaghers Syndrome
	(D)	Primordial prevention		(D)	Plummer-Vinson Syndrome
22.	Sho	e leather survey was carried out	27.	Soa	p bubble appearance is seen in :
	by:	o location out toy the out how out	•	(A)	Osteosarcoma
	(A)	Churchill H. V.		(B)	Osteoclastoma
	(B)	Frederick McKay		(C)	Ewing's sarcoma
	(C)	Trendley H. Dean		(D)	Multiple myeloma
	(D)	Greene Vardiman Black	28.		g used for the treatment of ostomia is:
23.	Exa	mple of a composite index is :	•	(A)	Atropine
	(A)	CPITN		(B)	Ephidrine
	(B)	DMF		(C)	Scopolamine
	(C)	TNM		(D)	None of the above
	(D)	Periodontal	29.	-	reflex is lost due to the paralysis
24.	Occ	urrence of disease clearly in	•	of:	th
	exce	excess of expected occurrence:		(A)	V <sup>th</sup> nerve
	(A)	Endemic		(B)	VII <sup>th</sup> nerve
	(B)	Pandemic		(C)	IX <sup>th</sup> nerve XII <sup>th</sup> nerve
	(C)	Sporadic		(D)	
	(D)	Epidemic	30.		n's syndrome is trisomy of:
25.	\//hi/	ch of the following is a measure		(A)	16 chromosome
<b>2</b>		spersion?		(B)	18 chromosome
	(A)	Mean		(C)	21 chromosome
	(B)	Median		(D)	23 chromosome
	` ,		31.		odontics treatment triangle was
	(C)	Mode  Other dend deviation			n by :
	(D)	Standard deviation		(A)	G. Z. Wright
ID	1D/3	1	(4)		Contd

	(B)	Henry Kempe	36.	Peri	od of mixed dentition growth
	(C)	Evangeline Jordan		spur	t in boys :
	(D)	Addleston		(A)	10-11 years
32.	Anto	rior bite plane is used for the		(B)	7-10 years
3Z.		ection of:		(C)	12-13 years
		据《····································		(D)	8-11 years
	(A)	Deep bite	37.	Whi	ch of the following GIC is using
	(B)	Crowding		as P	it and Fissure Sealent?
	(C)	Anterior cross bite		(A)	Туре 6
	(D)	Posterior cross bite	•	(B)	Type 8
33.	The	Daily dietary recommendation of		(C)	Type 7
1. 15	Calo	ium in children is :		(D)	Type 9
	(A)	600-800 mg	38.	The	first evidence of calcification of
	(B)	800-1200 mg		Prim	nary teeth begins approximately
	(C)	1200-1600 mg		at:	
	(D)	1600-2000 mg		(A)	11 weeks in utero
0.4		82. 1 The Control of		(B)	14 months in utero
34.		el content in Stainless steel		(C)	16 months in utero
		yn is :	-	(D)	6 months in utero
	(A)	17-19%	39.	Cla	ssical Conditioning was
	(B)	10-13%		prop	posed by:
	(C)	12-14%		(A)	Skinner
	(D)	11-12%		(B)	Pavlov
35.	Epsi	tein Pearls are found :		(C)	Peaget
	(A)	On crest of ridges		(D)	Hassel
	(B)	Along mid palatine raphe	40.	Inci	sal Liability is :
	(C)	Midline of mandible		(A)	Maxilla-3mm, Mand-5mm
	(D)	On the buccal surface of incisor		(B)	Maxilla-7mm, Mand-5mm
IB –	1D/3	4	(5)		(Turn over)

	(C)	Maxilla-5mm, Mand-7mm	45.	Ante	erior bolton's ratio is :	
	(D)	Maxilla-5mm, Mand-3mm		(A)	77.2%	
41.	디	Angle introduced his classi-		(B)	91.3%	
<b>-</b> † 1.		ion of malocclusion in :	•	(C <sub>i</sub> )	88.2%	
	(A)	1988		(D)	91.7%	
	(A) (B)	1899	46.	Mus	scle affected in torticollis :	
÷	• •			(A)	Lateral pterygoid	
	(C)	1901		(B)	Sternocleidomastoid	
	(D)	1801		(C)	Buccinators	
42.	The	most important synchondrosis for	•	(D)	Temporalis	
	the	growth modification is:	47.	Mid	line shift with unilateral posterior	
	(A)	A) Spheno ethmoidal			cross bite may be due to a change	
	(B)	Spheno occipital		in:		
	(C)	Inter sphenoidal		(A)	Pitch	
	(D)	Inter maxillary		(B)	Roll	
43.	Mon	nent of a force is :		(C)	Yaw	
	(A)	Force × perpendicular from		(D)	None of the above	
	( )	centre of rotation	48.	Ten	th theorem of retention was put	
	(B)	Force × perpendicular from		forth by:		
	(=)	centre of resistance		(A)	R. Riedel	
	(C)	Area under the stress strain		(B)	W. R. Proffit	
	(0)	graph		(C)	A. Lundstrom	
	(D)	Force × range		(D)	R. E. Moyers	
	` .		49.	Nor	mal naso labial angle necessi-	
44.		d order bends are :		tates	s the extraction of :	
	(A)	in-out bends		(A)	Upper first premolar	
	(B)	Tip back bends	•	(B)	Upper second premolar	
	(C)	Torquing bends		(C)	Upper incisor	
	(D)	Anti-rotation bends		(D)	None of the above	
IB-	1D/34	1	(6)		Contd.	

50.	Reve	erse orthodontics refers to :		(C)	Square
	(A)	Repeated orthodontic treat- ment		(D)	Rhomboid
	(B)	Camouflage treatment	55.		ch of the following is not a groove
	(C)	De-compensation		•	tern of mandibular second
	(D)	Treatment using lingual		•	nolar?
		appliances		(A)	Y
ΕĄ	Mhi	ch of the following deciduous		(B)	H
51.		h does not resemble any of the		(C)	U
		n in permanent dentition?		(D)	M
	(A)	Α	56.	Cell	s concerned with resorption of
	(B)	В		dec	iduous teeth:
	(C)	J		(A)	Osteoblast
	(D)	L		(B)	Cementoclast
52.	₩hi	ch of the following permanent		(C)	Odontoclast
<b>UL</b> .		h has a single antagonist in		(D)	Pericytes
	occl	usion?	57.	Whi	ich of the following tooth contain
	(A)	Maxillary third molar		sup	plementary grooves?
	(B)	Mandibular third molar		(A)	Maxillary second premolar
	(C)	Mandibular canine		(B)	Mandibular second molar
	(D)	Maxillary central incisor		(C)	Both (A) and (B)
53.	Wh	ich of the following tooth has		(D)	None of the above
		sial cusp slope greater than distal	58.	Col	is seen in :
	cus	p slope ?	50.	(A)	Buccal mucosa
	(A)	Maxillary first premolar			•
	(B)	Permanent mandibular canine		(B) <sub>.</sub>	
٠	(C)			(C)	
	(D)	Maxillary second premolar		(D)	Hard palate
54.		pe of occlusal aspect of maxillary	59.	•	thelial rests of Malassez is seen
	peri	manent first molar is :		in:	
	(A)	Trapezoidal		(A)	Palate
	(B)	Rectangle		(B)	Gingiva
IB-	-1D/3	34	(7)		(Turn over)

Mucoepidermoid carcinoma (C) Periodontal ligament (C) Dental pulp (D) Sebaceous carcinoma (D) Mesiolingual groove is seen in which 65. Odontoclastoma is: of the following teeth? (A) Giant cell tumor of bone Maxillary lateral incisor (A) (B) Giant cell granuloma (B) Mandibular first premolar (C) Internal resorption of tooth (C) Permanent mandibular first (D) Hypercementosis molar 66. Mumps is caused by: Mandibular canine (D) Orthomyxovirus (A) Leong's Premolar is: 61. Adenovirus (B) Dens invaginatus (A) (C) Paramyxovirus (B) Dens evaginatus (D) Coxsackie virus (C) Taurodontism 67. Odontoclasia is: Enamel hypoplasia (A) Linear enamel caries Most common benign tumor of oral (B) Tooth resorption cavity is: (C) Odontogenic tumor (A) **Papilloma** (D) Dry socket (B) Nevus 68. Bifid rib is seen in which of the (C) Fibroma following syndrome? (D) Lipoma (A) Papillon Leferve syndrome 63. Ackerman's tumor is: (B) Gorlin syndrome Pleomorphic adenoma (A) Sturge Weber syndrome (C) (B) Verrucous carcinoma (D) Ascher syndrome Spindle cell carcinoma (C) Most common oral manifestation of 69. (D) Adenolymphoma diabetes mellitus is: 64. Which of the following salivary gland (A) Leukoplakia tumor is mimic mucocele? (B) Periodontal abscess

ANUG

Hyperplastic candidiasis

Contd.

(C)

(D)

(8)

Adenoid cystic carcinoma

Salivary duct carcinoma

(A)

(B)

IB - 1D/34

60.

	(A)	Trigeminal neuralgia		mar	ndible is :
	(B)	Temporal arteritis		(A)	Lateral Pterygoid
	(C)	Sphenopalatine neuralgia		(B)	Medial Pterygoid
	(D)	Glossopharyngeal neuralgia		(C)	Temporalis
	(1)	Olossopharyngearnediaigia		(D)	Masseter
71.	Drug	of choice for the treatment of	76.	June	ctional epithelium shows rete peg
	Actir	nomycosis is :		form	nation in :
	(A)	Penicillin		(A)	Stage I gingivitis
	(B)	Cephalosporin		(B)	Stage II gingivitis
	(C)	Clotrimazole		(C)	Stage III gingivitis
	(D)	Metronidazole		(D)	Stage IV gingivitis
72.	Мое	ller's Glossities is seen in :	77.		nion of connective with root ace a root surface that has been
	(A)	Iron deficiency Anaemia		path	ologically exposed :
	(B)	Vitamin B6 deficiency		(A)	Reattachment
	(C)	Vitamin B12 deficiency		(B)	New attachment
	(D)	Thiamin deficiency		(C)	Regeneration
70				(D)	Repair
73.		span of platelet is :	78.	lf ha	If of the crown along with root is
	(A)	6-8 days		=	arated or one half is removed, it
	(B)	12-14 days		is ca	alled:
	(C)	60 days		(A)	Radisection
	(D)	120 days		(B)	Hemisection
74.	Rusk	nton bodies are present in :		(C)	Apexification
r -T.		•		(D)	Coronosection
	(A)	Periodontal cyst	79.		t abundant immunoglobulin in the
. *	(B)	Dentigerous cyst		seru	ım is :
	(C)	Odontogenic keratocyst		(A)	lgG
	(D)	Stafne's cyst		(B)	IgA
IB≟∘	1D/34	<b>!</b>	(9)		(Turn over)

70. Alarm clock headache is:

75. Muscle which depresses the

	(C)	lgM		(C)	4 months
	(D)	lgE		(D)	2 months
80.		ch of the following grafts has the	85.		microorganism found in higher
	best	t prognosis?		titer	s in pregnancy gingivitis :
	(A)	Xenograft		(A)	P. gingivalis
	(B)	Autograft		(B)	P. intermedia
	(C)	Allograft		(C)	Capnocytophaga
	(D)	Homograft		(D)	A. viscosus
81.	Ran	nping can be done for a :	86.	Eac	h ml of 2% lignocaine contains :
	(A)	One wall defect		(A)	0.2 mg of lignocaine
	(B)	Two wall defect		(B)	2 mg of lignocaine
	(C)	Three wall defect		(C)	10 mg of lignocaine
	(D)	Zero wall defect		(D)	20 mg of lignocaine
82.	Halir	meters are used in :	87.		ch of the following mandibular
	(A)	Oral malodour			h may be removed without pain
	(B)	Tooth mobility			the administration of IANB and
	(C)	Pathologic migration		_	ıal block?
	(D)	Pulp testing		(A)	All molars
83.	Can	ant of full mouth disinfaction		(B)	Both premolars and molars
00,		cept of full mouth disinfection put forward by:		(C)	All teeth in lower jaw
	(A)	Miller		(D)	First premolar and canine
	(B)	Glickman	88.	Follo	owing structures are present in
	(C)	William		the s	ubmandibular space except:
	(D)	Quirynen		(A)	Deep part of submandibular
		•			gland
84.	-	nelisation of gingivectomy		(B)	Facial artery
	wour	nd is completed by:		(C)	Mylohyoid nerve and vessels
	(A)	4 days		(D)	Superficial part of sub-
	(B)	1 month			mandibular gland
IB – ′	1D/34	(	10)		Contd.

- 89. Surgical emphysema may develop immediately after:
  (A) Cutting the tooth with airotor
  (B) Cutting the tooth with micromotor
  (C) Cutting the bone with micromotor
  (D) Cutting the bone with airotor
- 90. Metronidazole is a :
  - (A) Quinolone
  - (B) Anti-fungal
  - (C) Imidazole derivative
    - (D) Antiviral
- 91. Which of the following artery does not accompany the nerve throughout its course?
  - (A) Lingual
  - (B) Infra orbital
  - (C) Inferior alveolar
  - (D) Posterior superior alveolar
- 92. Secondary bony callus formation takes approximately:
  - (A) 1-2 weeks
  - (B) 3-4 weeks
  - (C) 4-6 weeks
  - (D) 6-8 weeks
- 93. Which of the following is DANGER SPACE?
  - (A) Pterygo mandibular
  - (B) Lateral pharyngeal

- (C) Superficial and deep temporal
- (D) Space between alar and prevertebral fascia
- 94. Weber Fergussion incision is used for:
  - (A) Mandibulectomy
  - (B) Maxillectomy
  - (C) Condylectomy
  - (D) Rhinoplasty
- 95. Cranial nerve passing through the centre of the cavernous sinus is:
  - (A) Optic nerve
  - (B) Ophthalmic nerve
  - (C) Occulo motor nerve
  - (D) Abducent nerve
- 96. Articular disc of TMJ is:
  - (A) Fibro cartilage
  - (B) Hyaline cartilage
  - (C) Elastic tissue
  - (D) TMJ ligament
- 97. Four clotting factors synthesized in liver are:
  - (A) III, V, VII and VIII
  - (B) II, VII, IX and X
  - (C) II, VI, IX and XII
  - (D) II, V, VII and IX
- 98. Which of the following is efficacious modality in diagnosis of salivary gland neoplasm?
  - (A) Incisional biopsy

- (B) Excision biopsy
- (C) Fine needle aspiration cytology
- (D) MRI scan
- 99. 1:1000 adrenaline ampule contains:
  - (A) 0.1 mg/ml
  - (B) 1 mg/ml
  - (C) 10 mg/ml
  - (D) 0.01 mg/ml
- 100. The target of gow gates technique is:
  - (A) Coronoid notch
  - (B) Sigmoid notch
  - (C) Mandibular foramen
  - (D) Neck of condyle
- 101. Calcifying epithelial odontogenic tumour commonly occurs in:
  - (A) Mandibular Posterior area
  - (B) Maxillary Posterior area
  - (C) Mandibular Anterior area
  - (D) Maxillary Anterior area
- 102. The most common histological variant of Ameloblastoma is:
  - (A) Follicular
  - (B) Plexiform
  - (C) Acanthomatous
  - (D) Granular cell
- 103. The histological variant of ameloblastoma with highest recurrenence rate is:
  - (A) Follicular
  - (B) Acanthomatous

- (C) Granular cell
- (D) Basal cell
- 104. Smoking is associated with the genesis of which of the salivary gland tumor?
  - (A) Pleomorphic adenoma
  - (B) Warthin's tumor
  - (C) Necrotizing sialometaplasia
  - (D) Papillary cystadenoma
- 105. A tumor which arises in salivary gland tissue and entrapped in parotid lymph nodes during embryogenesis is:
  - (A) Pleomorphic adenoma
  - (B) Warthin's tumor
  - (C) Oncocytoma
  - (D) Mucinous adenocarcinoma
- 106. In which of the following condition the normal flora becomes pathogenic?
  - (A) Diabetes
  - (B) Leukemia
  - (C) AIDS
  - (D) All of the above
- 107. Following are the beneficial effects of the resident flora except:
  - (A) Synthesis of Vitamin K
  - (B) Production of Pencillinase
  - (C) Suppression of colonization by pathogen
  - (D) Bacteriocin production

108.	Which of the following can be congenital infection?		a :	113.	Plasmid mediated resistance was first detected in :		
	(A)	AIDS			(A)	Salmonella	
	(B)	Toxoplasmosis			(B)	Shigella	
	(C)	CMV		-	(C)	Proteus	
	(D)	All of the above			(D)	E. coli	
109.		owing bacteria produce rotoxin except :	s '	114.	exce	•	
	(A)	Staph aureus			` ,	Rabies	
	(B)	B. anthracis			(B)	Toxoplasmosis	
	(C)	B. cereus			(C)	Hydatid	
	(D)	V. cholerae			(D)	Tetanus	
110.		ch of the following disease is no ired by ingestion ?	ot	115.	*	I mediated transmission of etic material in bacteria is known	
	(A)	Salmonellosis			(A)	Transcription	
	(B)	Shigellosis			(B)	Translation	
	(C)	Diptheria			(C)	Conjugation	
	(D)	Cholerae			(D)	Transduction	
111.	Follo exce	owing bacteria are invasive pt:	∌ ,	116.	Which is the cement with the highest solubility?		
	(A)	Shigella			(A)	Ca(OH) <sub>2</sub>	
	(B)	S. Typhi			(B)	Silicate	
	(C)	Vibiro Cholerae			(C)	Zn(PO <sub>4</sub> )	
	(D)	N. Meningitides			(D)	GIC	
112.		h of the following is vector borne	9 1	117.	Adva	intage of LED ligth is :	
	disea	ase?			(A)	Small and portable	
	(A)	Typhus			(B)	Low power consumption	
	(B)	Dengue			(C)	Minimal deterioration of light	
	(C)	Yellow fever				output	
	(D)	All of the above			(D)	All of the above	
IB – 1	ID/34		(13)			(Turn over)	

118.	8. The angle between rake face and		123.	Cement which is kindest to dental		
	radia	al line is :		pulp	:	
	(A)	Rake angle		(A)	GIC	
	(B)	Clearance angle		(B)	Polycarboxylate	
	(C)	Blade angle		(C)	Silicate	
	(D)	Radial clearance		(D)	Resin cement	
119.	Eam	es technique provides for :	124.	Corr	rugated gold is a type of	
	(A)	Low mercury-alloy ratio		(A)	Casting alloy	
	(B)	High mercury-alloy ratio		(B)	DFG	
	(C)	Easy waste management		(C)	Technic alloy	
	(D)	Good polishing		(D)	None of the above	
120.	Whi	ch of the following is an	125.	Elas	stic modulus of a material	
	estro	ogenic material?		indic	cates its:	
	(A)	Bisphenol – A		(A)	Rigidity	
	(B)	Poly acrylic acid		(B)	Flexibility	
	(C)	Nickel		(C)	Proportional limit	
	(D)	DFG		(D)	Viscosity	
121.	Simu	ultaneous application of acid to	126.	The	weakest part of mandible where	
	enar	nel and dentine is known as :		fract	ure occurs is :	
	(A)	Total etch technique		(A)	Neck of condyle	
	(B)	Dry bonding		(B)	Angle of mandible	
	(C)	Wet bonding		(C)	Midline	
	(D)	Hybridization		(D)	Canine fossa	
122.	Sme	ar layer consists of :	127.	Cleft	palate repair is ideally done at :	
	(A)	Enamel debris		(A)	Birth	
•	(B)	Microorganism		(B)	1-6 months	
	(C)	Dentin chips		(C)	7-18 months	
	(D)	All of the above		(D)	24-36 months	
IB-	1D/34	(	14)		Contd.	

128.		most commonly involved bon	e in	(C)	Polydioxone (PDS)
	frac	ture of the face is:		(D)	Polyglyconate (Chromic catgut)
	(A)	Nasal	13	3. Vita	amin K is required by liver for the
	(B)	Maxilla			duction of active :
	(C)	Mandible		(A)	Fibrinogen
	(D)	Zygoma		(B)	Platelet
129.	Car	cinoma of the lower alved	olar	(C)	Globulins
	ridge	e occurs around :		(D)	Prothrombin
	(A)	Incisor	134	4 Exc	citation of baroreceptors in
	(B)	Canine			eries reflexly causes :
•	(C)	Molar	•	(A)	Peripheral vasoconstriction
	(D)	Pre molar		(B)	Arterial pressure to decrease
130.	Salivary stones occurs most commonly in :		nst	(C)	Increase cardiac output
				(D)	Vasovagal syncope
	(A)	Parotid	138	5. Car	diac output is equal to the product
	(B)	Sublingual		of:	an equal of the
	(C)	Submandibular		(A)	Stroke volume × venous return
	(D)	Submaxillary		(B)	Stroke volume × heart rate
121	Bone	•		(C)	Venous return × heart rate
131.		es, stones moans and groa es to :	1115	(D)	Stroke volume × cardiac index
	(A)	Hyperparathyroidism	136	. Pre	-optic area of hypothalamus is
	(B)	Hyperthyroidism		con	cerned with :
	(C)	Hypoparathyroidism		(A)	Milk ejection
	• •	•		(B)	Uterine contractility
	(D)	Hypothyroidism		(C)	Temperature regulation
132.	Thes	suture that maintains strength	for	`(D)	Memory
• •	longe	est time is :	137	. REN	M sleep is associated with :
	(A)	A) Polyglycolic Acid (Dexon)		(A)	Normal heart rate
	(B)	Polyglactin (Vicryl)		(B)	Increased muscle tone
IB – ′	1D/34		(15)		(Turn over)

- (C) Decreased respiratory rate
- (D) Extreme inhibition of peripheral muscles
- 138. Sympathetic stimulation:
  - (A) Increases gastric motility
  - (B) Causes Pupillary dilation
  - (C) Decreases cardiac activity
  - (D) Vasodilation
- 139. Which of the following best describes the Broca area of speech?
  - (A) Sensory speech area
  - (B) Controls emotions
  - (C) Located in temporal bone
  - (D) Motor speech area
- 140. Intentional tremor occurs in the disease of :
  - (A) Cerebrum
  - (B) Basal ganglia
  - (C) Cerebellum
  - (D) Spinal cord
- 141. Pain sensation travels through the:
  - (A) Posterior column
  - (B) Lateral spinothalamic tract
  - (C) Corticospinal tract
  - (D) Vestibulospinal tract
- 142. ADH increases the permeability of distal tubule to:
  - (A) Proteins

- (B) Amino acids
- (C) Glucose
- (D) Water
- 143. Haemoprotein which binds to oxygen most strongly is:
  - (A) Adult haemoglobin
  - (B) Foetal haemoglobin
  - (C) Haemosiderin
  - (D) Myoglobin
- 144. Deficiency of which one of the following does not lead to anemia?
  - (A) Serum Iron
  - (B) Serum Copper
  - (C) Vitamin C
  - (D) Serum Zinc
- 145. Most of the enzymes secreted in the human gastrointestinal tract are :
  - (A) Oxidases
  - (B) Reductases
  - (C) Lyases
  - (D) Hydrolases
- 146. The primary function of the salivary amylase is:
  - (A) Hydrolysis of dietary starch
  - (B) Hydrolysis of bacterial cell wall
  - (C) Hydrolysis of glycogen
  - (D) Hydrolysis of pectin

- 147. Final common oxidative pathway which integrates oxidative products of fats, proteins and carbohydrates is:
  - (A) Gluconeogenesis
  - (B) Glycolysis
  - (C) Citric acid cycle
  - (D) Anaerobic glycolysis
- 148. Largest amount of energy is produced by which of the following?
  - (A) Oxidation of amino acids
  - (B) Beta oxidation of saturated fatty acids
  - (C) Beta oxidation of unsaturated fatty acid
  - (D) Anaerobic glycolysis
- 149. The substance that contributes maximally to the osmolality inside the cell is:
  - (A) Protein
  - (B) Phosphate
  - (C) Sodium
  - (D) Urea
- 150. The unique feature of Mitochondria is its:
  - (A) Actin
  - (B) Myosin
  - (C) Prothrombin
  - (D) DNA

- 151. Many substances are removed from the cell to outside by:
  - (A) Phagocytosis
  - (B) Pinocytosis
  - (C) Exocytosis
  - (D) Chemotaxis
- 152. Nerve fibre with slowest conduction is:
  - (A) A alpha
  - (B) A gamma
  - (C) Beta fiber
  - (D) C fibres
- 153. The superior meatus of the nose contains the opening of :
  - (A) Anterior ethmoidal air cells
  - (B) Middle ethmoidal air cells
  - (C) Posterior ethmoidal air cells
  - (D) Maxillary air sinus
- 154. The posterior auricular nerve is a branch of:
  - (A) Chorda tympani
  - (B) Facial
  - (C) Auriculo temporal
  - (D) Accessory nerve
- 155. The lateral wall of the infratemporal fossa is formed by :
  - (A) Lateral pterygoid plate
  - (B) Zygomatic arch
  - (C) Ramus of mandible
  - (D) Carotid sheath

- 156. The lymphatic from the tip of the tongue drains into:
  - (A) Submandibular lymph nodes
  - (B) Submental lymph nodes
  - (C) Jugulodigastric lymph nodes
  - (D) Jugulo omohyoid lymph nodes
- 157. Which of the following is a branch of maxillary nerve?
  - (A) Supratrochlear
  - (B) Zygomaticotemporal
  - (C) Infratrochlear
  - (D) External nasal
- 158. The taste sensation from the posterior one-third of the tongue is carried by:
  - (A) Chorda tympani
  - (B) Lingual nerve
  - (C) Hypoglossal nerve
  - (D) Glossopharyngeal nerve
- 159. Which of the following muscle assists in the elevation of mandible?
  - (A) Medial pterygoid
  - (B) Lateral pterygoid
  - (C) Digastric
  - (D) Genohyoid
- 160. The sensory root of the facial nerve is known as:
  - (A) Lingual nerve
  - (B) Chorda tympani
  - (C) Nervous intermedius
  - (D) Greater petrosal nerve

- 161. Drug causing hypocalcaemia is:
  - (A) Lithium
  - (B) Phenytoin
  - (C) Thiazides
  - (D) Vitamin A intoxication
- 162. All are side effects of steroids except:
  - (A) Myopathy
  - (B) Diarrhoea
  - (C) Peptic ulcer disease
  - (D) Osteoporosis
- 163. Epinephrine causes all except:
  - (A) Increase in blood glucose levels
  - (B) Increase blood lactate
  - (C) Increases free fatty acids
  - (D) Increases triglycerides
- 164. Which of the following is an ester linked local anesthetic?
  - (A) Lidocaine
  - (B) Bupivacaine
  - (C) Benzocaine
  - (D) Ropivacaine
- 165. Which of the following is heparin antagonist?
  - (A) Warfarin
  - (B) Vitamin K
  - (C) Coumarin
  - (D) Protamine sulphate

166.		te Asthma is treated with :	171.	-	otype in a patient with Turner
	(A)			•	Irome is :
	(B)	Salbutamol		(A)	45XO
	(C)	Digoxin		(B)	47XXY
	(D)	Frusemide		(C)	46XX
167.	Phys	sical dependence is produced		(D)	47XYY
	by:		172.	The	most common site of intra-oral
	(A)	Caffeine		nevi	is:
-	(B)	Amphetamine		(A)	Hard palate
	(C)	Cocaine		(B)	Buccal mucosa
	(D)	Pethidine		(C)	Gingiva
160	lotoli	erance is :		(D)	Lips
169.	(A) (B) (C) (D) End	Toxic effect of a drug in an individual at therapeutic doses It indicates high threshold of the individual to the action of a drug  Converse of the tolerance  All of the above  ocrinal side effects of hotropic drugs are:  Gynaecomastia  Galactorrhea  Both of the above	174.	malign(A) (B) (C) (D) The	most common potentially gnant state of oral mucosa is: Leukoplakia Erythroplakia Submucosal fibrosis Keratoplakia most common malignant plasm of the oral cavity is: Squamous cell carcinoma Basal cell carcinoma Malignant melanoma Sarcoma
	(D)	None of the above	175.	Une	rupted teeth is a common
	` '				re of :
	Ť	of choice for febrile seizures :		(A)	Dentigerous cyst
	(A)	Diazepam		(B)	Odontogenic keratocyst
	(B)	Phenytoin		(C)	Adenomatoid odontogenic
	(C)	Carbamate			tumour
	(D)	Valproate		(D)	All of the above
IB1	D/34	(	19)		(Turn over)

- 176. Swan-neck deformity of the fingers is seen in :
  - (A) Osteoarthritis
  - (B) Rheumatoid arthritis
  - (C) Osteomalacia
  - (D) Osteoporosis
- 177. Arthus type allergic reaction is a feature of :
  - (A) Type I reactions
  - (B) Type II reactions
  - (C) Type III reactions
  - (D) Type IV reactions
- 178. Progression to chronic liver disease is seen most commonly with infection by:
  - (A) Hepatitis A
  - (B) Hepatitis B
  - (C) Hepatitis C
  - (D) Hepatitis D
- 179. Blood Coagulation Factor which is not required in the intrinsic pathway for coagulation is:
  - (A) Factor 9
  - (B) Factor 11
  - (C) Factor 7
  - (D) Factor 13
- 180. All the following are suggestive of iron deficiency anemia except:
  - (A) Koilonychia
  - (B) Low reticulocyte response

- (C) Decreased serum ferritin
- (D) Decreased Total Iron-binding
  Capacity (TIBC)
- 181. All are side effects of corticosteroids except:
  - (A) Myopathy
  - (B) Diarrhoea
  - (C) Peptic ulcer disease
  - (D) Osteoporosis
- 182. Vitamin B12 deficiency may cause all of the following except:
  - (A) Optic neuritis
  - (B) Myelopathy
  - (C) Myopathy
  - (D) Neuropathy
- 183. Which of the following is the most common site for a fracture associated with osteoporosis?
  - (A) Femur
  - (B) Hip
  - (C) Radius
  - (D) Vertebra
- 184. All are the causes of bilateral parotid enlargement except:
  - (A) Diabetes mellitus
  - (B) Systemic lupus erythromatosus
  - (C) HIV
  - (D) Sarcoidosis

185.		h of the following is not a clinical			(B)	Anti HBsAg
	featu	re of cerebellar disease ?			(C)	HBeAg
	(A)	Nystagmus			(D)	Anti HBeAg
	(B)	Hyperreflexia	1:	90.	Norr	nal Serum Calcium, Phosphate
	(C)	Hypotonia			and.	ALP are seen in :
	(D)	Tremors			(A)	Osteomalacia
186.		he following are Vitamin K-			(B)	Osteopetrosis
	exce	endent coagulation factors pt:			(C)	Osteoporosis
	(A)	Factor X			(D)	Osteonecrosis
	(B)	Factor IX	1	01	Nac	opharyngeal carcinoma mostly
	(C)	Factor III		<b>Ο</b> 1.	arises from the:	
	(D)	Factor VII			(A)	Roof of nasopharynx
187.	Excessive weight gain, menorrhagia,				(B)	Posterior wall
	cold intolerance and weakness are				(C)	Anterior wall
	suggestive of:				(D)	Lateral wall
	(A)	Obesity				
	(B)	Hypothyroidism	1	92.	_	hest incidence of naso-
	(C)	(C) Cushing Syndrome			phar	yngeal carcinoma is seen in :
	(D)	Anasarca			(A)	Indian
188.	Scre	ening test used for the diagnosis	i		(B)	Chinese
	of C	ushing syndrome is :			(C)	Americans
	(A)	ACTH stimulation test			(D)	Europeans
	(B)	Plasma Cortisol	1	93	Whi	ch region is most commonly
	(C)	Plasma FSH	•	00.		cted in actinomycosis?
	(D)	Plasma Aldosterone			(A)	Thorax
189.	Whi	ch serological marker is	•		(B)	Liver
	suggestive of highly infectious stage		•		` '	
	of ac	cute Hepatitis B?			(C)	Facio-cervical
	(A)	HBsAg			(D)	Liver
IB-	1D/34	1	(21)			(Turn over)

194. A punched out edge is characteristic (B) 6-9 weeks of which ulcer? (C) 10-24 weeks (A) Tuberculosis (D) 25-36 weeks (B) Malignant 198. Dentigerous cyst develops around: (C) Rodent (A) Root (D) **Syphilis** (B) Crown 195. Cystic hygroma is a: (C) Gum (A) Dermoid cyst (D) Gingival Margin (B) Cavernous haemangioma 199. In tongue cancers, the least affected (C) Lymphangiectasia site is: Sebaceous cyst (A) Ventral margin 196. The most common indication for the (B) Dorsal margin removal of sublingual salivary gland (C) Tip is: (D) Lateral margin (A) Ranula (B) Lymphangioma 200. Ameloblastoma most commonly (C) Neoplasm develops around: (D) Stone Incisors (A) **Premolars** (B) 197. Cleft lip is ideally repaired at what age? (C) Canine

(A) Soon after birth

(D)

Third molar

## SPACE FOR ROUGH WORK

